



Monitoring knowledge, risk perceptions, preventive behaviours and trust to inform pandemic outbreak response

29 July 2020

# SURVEY TOOL AND GUIDANCE

**Rapid, simple, flexible behavioural  
insights on COVID-19**

Address requests about publications of the WHO Regional Office for Europe to:

Publications  
WHO Regional Office for Europe  
UN City, Marmorvej 51  
DK-2100 Copenhagen Ø, Denmark

Alternatively, complete an online request form for documentation, health information, or for permission to quote or translate, on the Regional Office website (<http://www.euro.who.int/pubrequest>).

© World Health Organization 2020

*All rights reserved. The Regional Office for Europe of the World Health Organization welcomes requests for permission to reproduce or translate its publications, in part or in full.*

*The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.*

*The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.*

*All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either express or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use. The views expressed by authors, editors, or expert groups do not necessarily represent the decisions or the stated policy of the World Health Organization.*

Photo: © WHO/Jonathan Schoeps

## 1. Introduction

**This document provides guidance to Member States in the WHO European Region that wish to conduct behavioural insights studies related to COVID-19.**

The COVID-19 pandemic outbreak is placing an overwhelming burden on health systems and authorities to respond with effective and appropriate interventions, policies and messages.

A poorly timed and managed **pandemic response or transition phase** can threaten the gains collectively achieved. The pandemic and its restrictions may have affected mental and physical well-being, social cohesion, economic stability as well as individual and community resilience and trust (1-5).

In this complex context, understanding how, why and the context in which humans and communities respond allows to

- 1) anticipate unwanted scenarios and initiate mitigating measures; and
- 2) implement pandemic response measures that are better informed, situated, accepted and thus more effective.

Population surveys can explore perceptions, acceptance of restrictions, mental and physical health, behaviours, information needs, misperceptions and more.

**WHO Regional Office for Europe and partners are offering Member States a tool to gain such insights which**

- 1) is evidence-informed;
- 2) can be regularly applied;
- 3) is flexible to adjust to the changing situation;
- 4) follows high ethical standards.

A few countries have rapidly instigated studies to gain such insights, and more countries are urged to prioritize such efforts to inform and support other response measures.

The approach presented in this guidance document was developed based on a framework initiated by the University of Erfurt, Germany, German national health authorities and others (Box 3).

This guidance document introduces:

- guidance on the recommended process and steps
- a sample methodology
- advice for obtaining ethical clearance
- a suggested sample questionnaire (Annex 1)
- codes for data analysis and establishing a protected website for presentation of findings (Annex 2).

WHO Europe's Insights Unit and Health Emergencies Programme are offering support to countries for implementation. We urge all users of the tool to let us know their plans so that we can coordinate and share. This way, we can prevent that two institutions in one country are working in parallel on the same type of study.

Please contact Katrine Bach Habersaat ([habersaatk@who.int](mailto:habersaatk@who.int)) or Martha Scherzer ([scherzerm@who.int](mailto:scherzerm@who.int)).

### **Box 1: Suggested approach to behavioural insights research for COVID-19**

Any country interested may use the questionnaire (Annex 1) to collect data regarding the public's risk perceptions, behaviours, trust, knowledge and other variables. This data is collected via online panels, and a national research group or private company can be engaged to collect the data. Using a set of codes (Annex 2) the findings can be automatically and immediately transferred to a protected webpage, if desired (examples of what this looks like in Figure 1). The webpage can be easily developed as a subpage on an existing website as decided by the implementing country. This allows national pandemic response groups to use the findings to inform pandemic response measures. In more detail:

- It is suggested to conduct a serial, cross-sectional study: Data can be collected repeatedly, e.g. weekly, with different participants so the status quo can be assessed over time.
- Repeated assessment of the same core variables allows analyzing changes over time. It also allows comparisons across countries if desired.
- Changing some variables allows adaptive research to a dynamic situation (e.g. asking for the acceptance of hand washing may be more important in the beginning while adding acceptance of closing schools or a lockdown may be relevant later in the epidemic).
- This allows in-time and adaptive monitoring of the variables (such as risk perceptions, knowledge, trust, behaviours and more) – and identifying changes over time to assess the relations between them.
- Variables can be adapted to different countries, target groups, cultural contexts and to the evolving situation and epidemiology over time. Randomization of answer options can be made where suitable.
- An automated data analysis website ensures immediate data analysis and provides fast access to the results (commented code for data analysis and website in Annex 2).
- Changes in risk perceptions or knowledge can be assessed over time.
- Data on acceptance of new response measures can be made rapidly available.
- Emerging issues, e.g. related to stigma can be identified as they emerge.
- National teams using the tool are urged to work in partner coalitions to discuss insights gained and implications for outbreak response interventions, policies and messages.
- Results can be made available to the media to support high quality and responsible reporting.

### **Box 2: Who is involved?**

- National or subnational health authorities will lead the study in their country.
- Research institutions may be engaged to conduct or support the study by health authorities. They may also take the initiative, approach health authorities and suggest collaboration.
- WHO Regional Office for Europe developed this guide and are offering support to European Region Member States for coordination or implementation.
- University of Erfurt, Germany originally developed the guidance and questionnaire and supported the adaptation made in this document. All documents were made available at no cost.

### **Box 3: Acknowledgements**

The following researchers and authorities were involved in developing this guidance

- Universität Erfurt (Cornelia Betsch (PI), Lars Korn, Lisa Felgendreff, Sarah Eitze, Philipp Schmid, Philipp Sprengholz)
- Robert Koch Institut (Lothar Wieler, Patrick Schmich)
- Leibniz Institute for Psychology Information (Michael Bosnjak)
- Bernhard Nocht Institute for Tropical Medicine (Michael Ramharter)
- Science Media Center (Volker Stollorz)
- Yale Institute for Global Health (Saad Omer)

## 2. Process: getting started

Figure 2 describes the suggested process of adapting the tool presented in this guidance document.

For steps marked with (\*) in Figure 2, standard materials can be found in this guidance document:

- standard guidance for adaptation
- questionnaire
- commented codes for data analysis website (Annex 2)


### Key steps include:

- Reaching out to WHO Regional Office for Europe for coordination and possible support ([habersaatk@who.int](mailto:habersaatk@who.int) and [scherzerm@who.int](mailto:scherzerm@who.int)).
- Liaising with key partners in the country. If you are a researcher: notifying country health and pandemic response authorities of the interest in using this tool and suggesting collaboration.
- Using current document to prepare country action plan with timeline plus roles and responsibilities
- Considering if the use of the tool will impede any national emergency response efforts. If so, please contact the WHO Regional Office for Europe before proceeding for guidance and support
- Deciding on which stakeholders to involve in the planning and which stakeholders should get access to the data.
- Adapting the guidance and questionnaire to the national context, consulting with the local community and key stakeholders to ensure clarity, applicability and cultural sensitivity (Annex 1).
- Seeking ethical clearance.
- Deciding on the data collection mechanism (online, Computer-assisted telephone interviewing (CATI), Computer-assisted personal interviewing (CAPI), mixed-mode).
- Deciding on the frequency of data collection. This decision should be made by country-level authorities based on criteria including the phase of the pandemic, ability of authorities to incorporate data and adjust response accordingly and the available human resources.
- Deciding on the sample. A cross-sectional sample is suggested (e.g. representative for age, gender, district – with a minimum of N = 1,000). A final sample size may be adjusted in consultation with the data collecting agency to be appropriate for the country.
- Entering an agreement with a data collector, e.g. a national statistics agency, an academic institution or a market research agency.
- If desired and appropriate, posting the protocol and questionnaire on PsychArchives.org: <https://hdl.handle.net/20.500.12034/2392>
- Collecting data: surveys are generally made available online for 38-48 hours. Ideally, any possible telephone-assisted data collection would happen in the same timeframe.
- Setting up an automated data analysis website for presentation of the data (using any website preferred and the codes provided in Annex 2). This website can use the open source R statistical package to analyse data and produce visual representations that can be easily reviewed and understood by a wide audience.
- Sharing results with national authorities who acknowledged their buy-in at the beginning: once results are available, we suggest regular meetings between all partners to share findings and discuss implications for pandemic response. As appropriate, the WHO Country Office or Regional Office for Europe will be available to participate in these meetings, with the clear understanding that the data

and subsequent decisions belong to the country, to provide input on the interpretation of data and how this might inform national pandemic response. Discussion includes possible actions to be taken in response to results and changes being seen over time and planning for a possible new round of data collection.

Please note that for suggested methodology, ethical clearance, sampling and more, there is detailed guidance below.

**Figure 2: Recommended steps and process**



Upload link: <http://dx.doi.org/10.23668/psycharchives.2782>

(\*) This guide provides supporting materials for this step (questionnaire, guidance, code).

### 3. Standard Approach

The following is the recommended standard approach for using this tool. The pandemic situation is evolving rapidly, and the most current version of the standard approach can be shared by WHO as an adaptable word version. Please contact Katrine Habersaat ([habersaatk@who.int](mailto:habersaatk@who.int)) or Martha Scherzer ([scherzerm@who.int](mailto:scherzerm@who.int)).

#### Flexibility and adaptation

As the COVID-19 pandemic evolves and the epidemiological and response situation rapidly changes, the study must be continuously updated so that the questions asked reflect the situation and provide the necessary information to shape effective and appropriate outbreak response measures and next steps.

National teams using the tool are encouraged to draw on the existing evidence-base from previous outbreaks and epidemics and ethical frameworks for decision making in public health (17,18) as findings emerge.

#### Aims and objectives of the study

The study will be initiated by health authorities in individual countries to gain insights into risk perception, knowledge, trusted sources of information, attitudes toward pandemic response initiatives and other variables **to inform COVID-19 outbreak response measures, including policies, interventions and communications.**

The primary objectives are to:

- Monitor variables that are critical for population behaviour to control transmission of the novel coronavirus, including risk perceptions, knowledge, self-efficacy, confidence in institutions, behaviours, rumours, affect, worry, resilience, trust in/use of information sources and more.
- Document changes over time in these factors to understand the effect of the pandemic process, new developments, events or measures taken.
- Monitor possible issues, e.g. related to misinformation or distrust, as they emerge, to allow early response.
- Identify relationships between variables to identify levers for effective and appropriate responses.
- Explore the relationship of psychological variables (e.g. worry, resilience, trust, affect) with the epidemiological situation and the events and measures taken.
- Identify gaps between perceived and actual knowledge.
- Evaluate the effectiveness of pandemic response measures, and the acceptance and effectiveness of policies and restrictions implemented, including the easing of such restrictions.

The secondary objectives are to:

- Contribute to post-outbreak evaluation, thereby contributing to the continued regional/global efforts to better understand mechanisms of crisis response.
- If additional research capacity is available, the data can be triangulated with data on media reporting, COVID-19 cases and other.



- If additional research capacity is available, the data can be triangulated with data on media reporting, imported or confirmed cases, etc.: The relationship between psychological variables and characteristics of the outbreak situation can be explored (i.e. how closely the perceived risk mirrors reported cases, relative import risk, media reports).

This approach allows a citizen-centred approach where insights into population perceptions and behaviours inform COVID-19 actions, alongside epidemiological data and considerations of economic, cultural, ethical, structural political nature and other.

### Study methods

The study method is decided upon by the national pandemic response authorities based on feasibility and appropriateness in the country. A 15-20 minutes online questionnaire in a serial cross-sectional design with multiple data collections is suggested as a standard approach (i.e., each sample will consist of different participants). Computer-assisted telephone interviews (CATI) surveys can be considered as additional or alternative data collection methods in countries where access to computers or smartphones is less widespread. For as long as social distancing is recommended, we do not recommend household surveys.

This is an observational study with voluntary participation in the general population, with expected low risk for participants. Potential risks identified include only the inconvenience of the time taken to respond to the survey, and given the current restrictions people face, many individuals currently have more available time. The variables and information requested does not allow to identify specific ethnic or disadvantaged population groups. Due to strict data protection measures, any risk related to non-anonymous publishing of data from the survey is considered very low, and the personal harm for the individual respondent related to such unlikely event is also considered low due to the less sensitive nature of the responses provided. Benefits include the sense of contributing and being able to participate in shaping the country's pandemic response.

It is suggested to collect data repeatedly (e.g. weekly or adapted to the epidemiological situation). This will allow to:

- identify developments over time (e.g. a decline in trust, or a decline in motivation to follow recommended behaviours);
- Identify new issues as they emerge (e.g. related to conspiracy theories, new misperceptions, stigma against certain groups or other) and address these;
- Detect effect or adverse responses to new restrictions, messages or actions taken.

Note that the cross-sectional design will not allow the assessment of actual causal relations and will only be snapshots of a current state of the public perceptions and behaviours.

In case of unexpected developments or new outbreak response measures implemented, pandemic response authorities may decide to change the time frame between the data collections.

If the survey is conducted repeatedly, it is advised to ensure a system to avoid participants to respond to the survey more than once. E.g. ensuring that only participants with a different ID can participate in

subsequent surveys. This procedure ensures both anonymity of participants and identification of duplicates.

A longitudinal panel would be an alternative to a serial cross-sectional design. This would mean asking the same participants repeatedly. It is not recommended to ask participants to fill in the same questionnaire every week, as answering the questions will potentially influence the answers given.

## Variables

Variables being surveyed include the following (see Table 1 for details):

- Socio-demography
- COVID-19 personal experience
- Health literacy
- COVID-19 risk perception:
  - Probability and Severity
  - Preparedness and Perceived self-efficacy
  - Prevention – own behaviours
- Affect
- Trust in sources of information
- Use of sources of information
- Frequency of Information
- Trust in institutions (perceptions)
- Policies, interventions (perceptions)
- Conspiracies (perceptions)
- Resilience (perceptions)
- Testing and tracing
- Fairness (perceptions)
- Lifting restrictions (pandemic transition phase)
- Unwanted behaviour
- Wellbeing
- COVID-19 vaccine

Variables include a combination of knowledge and behavioural questions that can only be answered by an individual based on the current situation along with other more complex constructs. Asking people to what degree they are following suggested prevention interventions such as hand washing and social distancing shows how many people self-report contributing to the response in these ways.

Other constructs are more complex and require validated questions to accurately assess, such as risk perception, self-efficacy, trust, affect, fairness, prevention, resilience, worry and conspiracy thinking. These variables are measured using validated questions or adapted validated questions. See **Table 1** under Methodology: Survey Tool below for details.

**Table 1.: Questionnaire – validation and value of variable and items included**

Variable	Items relate to	Validation of psychological construct	Value in relation to study objectives	Standard analysis conducted
<b>Socio-demography</b>	Age, gender, education, medical background, chronic illness, rural/urban, district, household, financial situation  ( <u>risk group</u> identified as: 70+ years and/or chronic illness)	Not a psychological construct	Allows stratifying findings per population groups	Summary provided at the end
<b>COVID-19 personal experience</b>	COVID-19 infection (own, someone close)	Not a psychological construct	Allows stratifying findings per infected (affected)/not infected (affected)	Summary provided at the end
<b>Health literacy</b>	Assessment of ease/difficulty in finding information on symptoms, finding out what to do if infected, understand what authorities say, judge reliability of information, follow recommendations, decide on prevention behaviours	Construct: health literacy  <u>Items adapted from:</u>  Sørensen K, Van den Broucke S, Pelikan JM, et al. Measuring health literacy in populations: illuminating the design and development process of the European Health Literacy Survey Questionnaire (HLS-EU-Q). BMC Public Health. 2013;13:948. Published 2013 Oct 10. doi:10.1186/1471-2458-13-948, and Griebler, Robert; Nitsche, Michael (2020): The Austrian Corona Health Literacy Questionnaire. Vienna: Gesundheit Österreich GmbH & Das Österreichische Gallup Institut	Allows to assess the individual perspective on access to, understanding of and use of information/knowledge	<u>Regression</u> is conducted for: <ul style="list-style-type: none"> <li>• What to do in case of suspected COVID-19</li> <li>• Decide when to engage in social activities</li> <li>• Understand restrictions and recommendations</li> </ul>
<b>COVID-19 risk perception: Probability and Severity</b>	Self-assessed probability and susceptibility to of contracting COVID-19	Psychological construct: risk perception.  <u>Validated items adapted from:</u>  Brewer, N. T., Chapman, G. B., Gibbons, F. X., Gerrard, M.,	Allows to identify possible patterns in behaviours/perceptions (see	<u>Regression</u> is conducted for: <ul style="list-style-type: none"> <li>• Probability</li> <li>• Susceptibility</li> <li>• Severity</li> </ul>

Variable	Items relate to	Validation of psychological construct	Value in relation to study objectives	Standard analysis conducted
	Self-assessed severity in case of contracting COVID-19	McCaul, K. D., & Weinstein, N. D. (2007). Meta-analysis of the relationship between risk perception and health behavior: the example of vaccination. <i>Health psychology, 26</i> (2), 136.	below) related to risk perceptions	
<b>Preparedness and Perceived self-efficacy</b>	Self-assessed COVID-19 self-protection and avoidance ability	<p>Psychological construct: preparedness</p> <p><u>Validated items adapted from:</u></p> <p>Bandura, A. (2006). Guide for constructing self-efficacy scales. <i>Self-efficacy beliefs of adolescents, 5</i>(1), 307-337.</p> <p>Psychological construct: perceived self-efficacy</p> <p><u>Validated items adapted from:</u></p> <p>Renner, B., &amp; Schwarzer, R. (2005). The motivation to eat a healthy diet: How intenders and nonintenders differ in terms of risk perception, outcome expectancies, self-efficacy, and nutrition behavior. <i>Polish Psychological Bulletin, 36</i>(1), 7-15.</p>	Allows to identify possible patterns in behaviours/perceptions (see below) related to self-efficacy	Results are shown for the entire sample
<b>Prevention – own behaviours</b>	Own behaviours: prevention measures (hand washing, avoid face, disinfectants, home when sick, physical distancing, face mask, antibiotics, not seeing family, friends)	<p>Psychological construct: prevention behaviour</p> <p><u>Items adapted from:</u></p> <p>Steel Fisher GK et al (2012). Public response to the 2009 influenza A H1N1 pandemic: a polling study in five countries. <i>Lancet Infectious Diseases 2012; 12: 845–50</i></p>	<p>Allows to compare knowledge and behaviour</p> <p>Allows to identify resiliency in upholding recommended behaviours which may need to be addressed</p>	<p>Results are shown for the entire sample.</p> <p>Knowledge of prevention and own prevention behaviours are compared for the entire sample (<u>see below</u>).</p> <p><u>Regression</u> analysis is conducted to identify characteristics of those who</p>

Variable	Items relate to	Validation of psychological construct	Value in relation to study objectives	Standard analysis conducted
				take up preventive behaviours  Specifically for <u>antibiotics</u> : please cross-reference with whether respondents had COVID-19, and whether it was confirmed by a test or not.
<b>Affect</b>	Affect related to COVID-19 (close, spreading, constant, fear-inducing, media hyped, helpless, stressful)	Psychological construct: affect  <u>Validated items adapted from:</u>  Bradley, M. M., & Lang, P. J. (1994). Measuring emotion: the self-assessment manikin and the semantic differential. <i>Journal of behavior therapy and experimental psychiatry</i> , 25(1), 49-59.	Allows to identify mental health implications of restriction – ultimately potentially as a warning sign that restrictions need to be changed	Results are shown for the entire sample  <u>Regression</u> is conducted for: <ul style="list-style-type: none"> <li>• Fear-inducing</li> <li>• Close-far away</li> <li>• Media-hyped</li> </ul>
<b>Trust in sources of information</b>	Trust in information sources (television, newspapers, health workers, social media, radio, Ministry of Health, Institute of Public Health, hotlines, official website, celebrities)	Psychological construct: trust  <u>Item ground in theory:</u>  Schweitzer, M. E., Hershey, J. C., & Bradlow, E. T. (2006). Promises and lies: Restoring violated trust. <i>Organizational behavior and human decision processes</i> , 101(1), 1-19.  Pearson, S. D., & Raeke, L. H. (2000). Patients' trust in physicians: many theories, few measures, and little data. <i>Journal of general internal medicine</i> , 15(7), 509-513.	Allows to identify trusted information sources, to be used for planning communications	Results are shown for the entire sample
<b>Use of sources of information</b>	Use of information sources (television, newspapers, health workers, social media, radio, Ministry of Health, Institute of Public Health,	Not a psychological construct	Allows to compare trust in and use of information sources and to identify widespread	Results are shown for the entire sample

Variable	Items relate to	Validation of psychological construct	Value in relation to study objectives	Standard analysis conducted
	hotlines, official website, celebrities)		sources, to be used for planning communications	
<b>Frequency of Information</b>	Frequency in information	Not a psychological construct	Allows to understand information needs, to be used for planning communications	Results are shown for the entire sample
<b>Trust in institutions (perceptions)</b>	Trust in ability of stakeholders to handle situation (Doctor, Employer, Hospitals, Ministry of Health, Institute of PH, Schools Universities, Kindergartens, Public transportation, Police, Church)	Psychological construct: trust  <u>Item grounded in theory:</u>  Schweitzer, M. E., Hershey, J. C., & Bradlow, E. T. (2006). Promises and lies: Restoring violated trust. <i>Organizational behavior and human decision processes</i> , 101(1), 1-19.  Pearson, S. D., & Raeke, L. H. (2000). Patients' trust in physicians: many theories, few measures, and little data. <i>Journal of general internal medicine</i> , 15(7), 509-513.	Allows to understand trust, and trends related to this, to be used for planning communications – and for detecting possible shifts in trust (e.g. following certain events or new restrictions) which can inform/promote/avoid future events.	Results are shown for the entire sample  <u>Stratification</u> is conducted for the following: <ul style="list-style-type: none"> <li>• Your own family doctor</li> <li>• Ministry of Health</li> <li>• Local Public Health Authority</li> </ul>
<b>Policies, interventions (perceptions)</b>	Perceptions related to possible/real government policies (COVID-19 vaccine, discrimination behaviours, testing, exaggeration in restrictions, quarantine)	Not a psychological construct	Allows to understand perceptions of policies, to inform policy planning, or to inform contingency plans before new policies.	Results are shown for the entire sample  Please include <u>numbers</u> for top 3 answers and top bottom answers to:  “If a vaccine becomes available and is recommended for me, I would get it” (Completely agree ++ vs. completely disagree ++)  <u>Regression</u> is conducted for: <ul style="list-style-type: none"> <li>• If a vaccine becomes available and is</li> </ul>

Variable	Items relate to	Validation of psychological construct	Value in relation to study objectives	Standard analysis conducted
				<p>recommended for me, I would get it.</p> <ul style="list-style-type: none"> <li>In the event of an outbreak it's appropriate to avoid certain people on the basis of their ethnicity.</li> <li>The government should be allowed to force people into self-isolation if they have been in contact with a person who was infected</li> <li>I think that the restrictions currently being implemented are greatly exaggerated</li> </ul>
<b>Conspiracies (perceptions)</b>	Perceptions related to transparency, motivations, monitoring, secrets, hidden organizations.	<p>Psychological construct: conspiracy thinking</p> <p><u>Validated items taken from:</u></p> <p>Bruder M, Haffke P, Neave N, Nouripanah N, Imhoff R. Measuring individual differences in generic beliefs in conspiracy theories across cultures: conspiracy mentality questionnaire. <i>Front Psychol.</i> 2013;4:225. Published 2013 Apr 30. doi:10.3389/fpsyg.2013.00225</p>	Allows to detect trends in possible conspiracy theories which may need to be addressed.	<p>Results are shown for the entire sample</p> <p><u>Regression</u> is conducted for psychological construct 'Conspiracy thinking' (questions combined)</p>
<b>Resilience (perceptions)</b>	Perceptions related to coping with stress and recovering. Ease/difficulty in not seeing friends and family.	<p>Psychological construct: resilience</p> <p><u>Validated items taken from:</u></p> <p>Smith, B. W., Dalen, J., Wiggins, K., Tooley, E., Christopher, P., &amp; Bernard, J. (2008). The brief resilience scale: assessing the ability to bounce back. <i>International journal of behavioral medicine</i>, 15(3), 194-200.</p>	Allows to identify mental health implications of restriction – ultimately potentially as a warning sign that restrictions need to be changed.	<p>Results are shown for the entire sample</p> <p><u>Regression</u> is conducted for psychological construct 'Resilience' (questions combined)</p>

Variable	Items relate to	Validation of psychological construct	Value in relation to study objectives	Standard analysis conducted
<b>Testing and tracing</b>	Barriers and drivers to getting tested and sharing names for tracing	Not a psychological construct. <u>Items grounded in theory:</u>  Michie et al (2014), The Behaviour Change Wheel. A Guide to Designing Interventions. Silverback Publishing. ISBN 978-1-912141-00-5.	Allows to understand barriers and drivers to testing and tracing	Results are shown for the entire sample  <u>Regression</u> is conducted for  <ul style="list-style-type: none"> <li>• I may not get tested</li> <li>• I may not share all names</li> </ul>
<b>Fairness (perceptions)</b>	Perceptions related to fairness of COVID-19 decisions (fair, would convince others)	Psychological construct: fairness  <u>Validated items taken from:</u>  Gamliel, E., & Peer, E. (2010). Attribute framing affects the perceived fairness of health care allocation principles. Judgment and Decision Making, 5(1), 11.	Allows to understand and possibly detect new trends in acceptance of restrictions which may inform new restrictions, lifting of restrictions, or the communication about these.	Results are shown for the entire sample  <u>Regression</u> is conducted for psychological construct 'Fairness' (questions combined)
<b>Lifting restrictions (pandemic transition phase)</b>	Perceptions related to lifting restrictions (adapted to country decisions made/considered)	Not a psychological construct	Allows to foresee reactions and perceptions concerning possible scenarios in the transition phase and to use these to inform decisions.	Results are shown for the entire sample  <u>Regression</u> is conducted for:  <ul style="list-style-type: none"> <li>• Selected statements as deemed relevant by country</li> </ul>
<b>Unwanted behaviour</b>	Reported own behaviour (discrimination, physical exercise, alcohol, diet, smoking, vaccination postponed, drugs against COVID-19, postponed doctor visit)	Not a psychological construct	Allows to identify adverse behaviours that may need to be addressed. Can be compared with data from doctors, supermarkets etc to assess validity of findings.	Results are shown for the entire sample  <u>Regression</u> is conducted for:  <ul style="list-style-type: none"> <li>• Exercised less than I usually do</li> <li>• Drank more alcohol than I usually do</li> <li>• Ate more unhealthy food than I usually do</li> <li>• Avoided going to the doctor with issues that</li> </ul>



Variable	Items relate to	Validation of psychological construct	Value in relation to study objectives	Standard analysis conducted
				could be postponed, e.g. vaccination or a check-up
<b>Well-being</b>	Rate of being cheerful, calm, active, fresh, day filled with interesting things	Psychological construct: Well-being  <u>Validated items from:</u>  WHO 5-item well-being scale (WHO-5). For systematic reviews, see Winther Topp et al 2015. Psychother Psychosom 2015;84:167–176, DOI: 10.1159/000376585.	Allows to identify mental health implications of restriction – ultimately potentially as a warning sign that restrictions need to be changed.	Results are shown for the entire sample  <u>Regression</u> is conducted:  • Across the five items
<b>COVID-19 vaccine</b>	Agreement with value statements re potential future vaccine (control COVID-19, help avoid restrictions, never accept it, should be mandatory).  Indication of own barriers/drivers to getting the vaccine (production country, recommendations, many vaccinated, free of charge, ease of access, used in other countries, COVID-19 risk, need if others are vaccinated)	Construct: Vaccine hesitancy  <u>Items grounded in theory:</u>  Michie et al (2014), The Behaviour Change Wheel. A Guide to Designing Interventions. Silverback Publishing. ISBN 978-1-912141-00-5 AND  Betsch, C., Schmid, P., Heinemeier, D., Korn, L., Holtmann, C., & Böhm, R. (2018). Beyond confidence: Development of a measure assessing the 5C psychological antecedents of vaccination. PLOS ONE, 13(12), e0208601. <a href="https://doi.org/10.1371/journal.pone.0208601">https://doi.org/10.1371/journal.pone.0208601</a> Open Data Open Materials	Allows to identify possible perceptions re a future COVID-19 vaccine as well as possible barriers and drivers, and to understand how these are related to demography and other variables.	Results are shown for the entire sample  <u>Regression</u> is conducted for:  • I believe a vaccine can help control the spread of COVID-19 • I would never accept to be vaccinated against COVID-19 • If a COVID-19 vaccine becomes available, it should be mandatory for all

### Data collection and analysis

Participants can be recruited via a trusted internal or external study sample provider as deemed feasible and appropriate by the national pandemic response authorities. The data collector may be:

- a trusted private market research agency
- an academic institution
- a government statistics agency
- other trusted data collection institution.

Users are recommended to share the current guidance and questionnaire with the data collector and discuss the process based on their available study tools and population panels. It needs to be specified in the agreement with data collectors that they need to follow data protection regulations as required by the state and regional authorities as well as GDPR, including as regards access to data, anonymity, protection of data, confidentiality and use of the data. The data must be collected in such a way where it will not be possible to identify individuals from the answers they have submitted.

It is recommended that a process of due diligence be conducted to evaluate the data practices of potential data collectors. Data collection and panel provider staff must have been provided with training on ethical considerations for the collection, use and storage of data (data security, data protection).

It is crucial to ensure the collected data is relevant, and not excessive in relation to the purpose for which it was obtained. The collected data should be exclusively used for the purpose of the current study and future related research. Users should ensure that the principles outlined in this document are met when working with data collectors.

Participants should take part in the survey voluntarily and can receive a remuneration, e.g. paid by the data collector. Remuneration should be agreed upon based on the usual procedures of the data collector and according to national standards. Each fielding period should be as short as possible as the situation evolves quickly, as do the peoples' perceptions. We suggest a maximum data collection period of maximum of 38-48 hours (e.g., 10am until 12pm the following day) with a maximum of two subsequent days.

After frequency of data collection (e.g. weekly) is decided on, each new data collection should take place with a new, independent sample.

The quota sample should match the current population in terms of age, gender and residency. Data collection can take place online or via phone (CATI).

Each country decides how they wish to analyse their data. One opportunity is offered with this tool: Based on a set of codes which have been pre-prepared for the questionnaire, the data (collected in an Excel file using specific labels) can be automatically and immediately displayed on a webpage which is protected by password. Each country decides which website is used for this (e.g. the ministry of health website, a website of a research institution or a website established just for this purpose). A design for the data display on this webpage has been created (in English). For a "dummy" presentation of how data is visualized, please see [https://projekte.uni-erfurt.de/cosmo2020\\_web/cosmo-analyses.html](https://projekte.uni-erfurt.de/cosmo2020_web/cosmo-analyses.html). (**Username:** web. **Password:** pWmG68qptP6AdhXLF4gZ9nQG8pNHQUSE). This automatic display of the findings allows for fast access to the results for multiple stakeholders (as decided by the implementers). More in-depth analysis of the data, e.g. stratifying of variables can be added at any time. Commented code for data analysis and website are available in annexes 2 and 3.

### Budget and timeline considerations

Costs associated with this study relate to data collection and human resources. Data collection may be conducted by government agencies as and where available. Some private sector market research firms have offered support for pro bono data collection, and this option can be explored on a country-by-country basis. Where third party, private data collection companies are hired, an estimated average cost per wave has been found to be \$2,000-\$4,000. The WHO Regional Office for Europe has some

emergency funding to support data collection as well as staff in WHO country offices and at the Europe Regional Office to provide support.

Initiation of data collection can begin quite quickly (potentially within one week) depending on coordination between government and researchers and availability of data collection platform.

### Institutional Review Board agreements, ethical standards met and safety monitoring

The study and handling of the data should follow all required regional and national data protection regulations. In general, data should be collected anonymously, with no collection of names, phone numbers, email addresses or other information which can identify participants or link participants to data. Only data from respondents aged 18+ will be included. If agencies collecting such data are hired, it is expected to hire only those agencies that have procedures to ensure this. If such data is collected it has to be anonymized before the data is analyzed.

Also, participants should provide informed consent before starting the questionnaire. Text on this is included in the questionnaire in Annex 1. The research contains negligible risks as there is no foreseeable risk of harm or discomfort other than potential inconvenience during participation. The study does not include deception and participants will be debriefed at the end of the survey. The study also involves only non-identifiable data about human beings.

### Ethical approval

Ethical approval should be sought at national level. Research is oriented on the ethical standards of American Association for Public Opinion Research (AAPOR) (<https://www.aapor.org/Standards-Ethics/AAPOR-Code-ofEthics.aspx>) and American Psychological Association (APA) (<https://www.apa.org/ethics/code/>).

Approval from the Ethical Review Committee of the WHO for the study and questionnaire may be sought on a country-by-country basis.

### Sample

Small effects may matter greatly on a population level, so a large sample size is recommended to allow for meaningful results. To obtain a high level of congruence between the distribution of the demographics in the sample and the adult population (regarding age, gender and living area), a sample size of  $n = 1000$  per wave is recommended.

Each data collection with  $n = 1000$  participants is suggested as a quota sample, matching the general population in the country in terms of age, gender and state/district.

The recommendations aim to decrease bias by retaining key proportions that are identical to those of the country population. Moreover, the goal of using a sample size of 1000 is to make the sample as representative as possible of the country population. The sample size of 1000 is a recommended number for surveys of large size populations. Research shows that the precision of estimates of surveys only increases very slightly beyond a sample size of 1000 (19). Thus, costs of inviting more than 1000 participants may exceed the statistical benefits.

## Tests

Analyses are suggested integrated in a R Notebook environment (for details, see Annex 2). All analyses are exploratory and may change based upon requirements of the situation. The data analysis script uses means of descriptive data presentation, regression analyses and correlation analyses.

Misinformation is collected as text fields and should be screened, summarized and offered to experts and those responsible for the crisis communication (e.g. to be debunked and inserted in FAQ lists).

Only completed data sets will be considered in the analysis. Missing values will be treated as missing values and not be imputed.

## Scientific review and validation of tools

The protocol and questionnaire were originally prepared by Professor Betsch at the University of Erfurt, Germany, and subsequently reviewed by a group of experts (Fig. 3), representing leading global experts in behavioural insights research for health and in developing and validating survey tools similar to the current. In addition, following each rounds of data collection in Germany, two scientists (Prof. Robert Böhm, University of Copenhagen, Denmark, and Britta Renner, University of Konstanz, Germany) have reviewed the data and how it was presented. To the degree possible, already validated items from previous surveys conducted were included in the questionnaire (e.g. questions related to risk perception, self-efficacy, trust, affect, fairness, prevention, resilience, worry and conspiracy thinking) (Table 1). The questionnaire as a whole has been validated through the six rounds of data collection in Germany which led to adjustments of the questionnaire.

It is recommended that in each country, the protocol and questionnaire are translated by an expert translator familiar with terminology of COVID-19 and behavioural science and with interview skills. It should then be reviewed by at least two national peer reviewers and revised accordingly. Reviewers should endorse the final protocol and questionnaire upon revision. The questionnaire should be pre-tested with a sample of respondents (age groups, gender, urban/rural) with a focus on their easy understanding of the questions before broad use.

## Limitations of the study

The urgency of the situation incurs some limitations to the study.

Using online panels limits the participation of certain important population groups, including the elderly (a risk group for COVID-19) and disadvantaged population groups such as migrants, refugees, young people below 18 years, homeless people and other vulnerable groups. Phone interviews as a supplement or instead of online panels can be used to mitigate this. Still, it may be assumed that some population groups will not take part in the survey, and so it cannot be claimed to represent their views, and the social benefit of the study may consequently be reduced. The findings of the survey need to be interpreted in this context. It may be considered to conduct supplementary more tailored and targeted surveys with specific population groups.

Since the findings related to the population at large may not apply to specific disadvantaged population groups, this affects the generalizability of the study findings. To overcome these limitations, health authorities are recommended to test recommended interventions informed by this survey with the broader population or specific population groups before rolling them out in a tailored fashion. This is

possible for e.g. specific messages or communication initiatives which can be tested with the target audience.

Due to the rapid development of the tool, it has not yet been possible to validate that, if the findings from these (weekly) surveys are used to inform pandemic response measures, this will directly lead to behaviour change in the public.

In addition, the complexity of the pandemic and crisis and the public response is considerable, and an online survey can only serve to monitor a few key issues - not explore them in-depth. Importantly, this survey can identify issues of concern that may need to be explored through other means, such as a supplementary qualitative telephone interview survey.

Another limitation of the study is that, while validated for other scales and well-grounded in robust behavioral research, the items have not been validated through a rigorous process for COVID-19 specifically. This is due only to the fact that we have never experienced this virus before and needs to be taken into account as a limitation in the interpretation of findings.

Self-reported behaviours are known to differ from actual behaviour, not least due to the social desirability effect, and so the findings related to behaviour should be interpreted with this reliability limitation in mind.

Finally, as each country adapts the questionnaire, not all data collected with this tool can be compared across countries for future evaluation purposes. The hope is that each country will collect and analyse at least several variables in common that may provide useful insights for cross-country comparison, but the main purpose of this tool is to help countries right now to determine the best approaches for their immediate COVID-19 response.

### Sharing the survey tool and guidance with other countries

If deemed appropriate, it is suggested that each user of the tool shares it via an open source research website to ensure methods and results can be shared with other countries.

## 4. Background: Review of relevant literature

Models of crisis and emergency risk communication (5) suggest that it is crucial to understand the risk perception of the population and the sources of information that they trust to enable effective communication and framing key messages. Messaging should be evidence-based and respond to misinformation and induce rational, adaptive and protective behaviour (6). However, little is known about the complex interplay of changing epidemiology, media attention, pandemic control measures, risk perception and public health behaviour (7). A study conducted during the influenza A(H1N1)pdm09 pandemic in 2009/2010 shows an “asynchronicity between media curves and epidemiological curves (...); media attention for influenza A H1N1 in Europe declined long before the epidemic reached its peak, and public risk perceptions and behaviours may have followed media logic, rather than epidemiological logic” (7). Thus, how people perceive the risk is not necessarily related to the actual risk. This perceived risk, nevertheless, influences protective behaviours (8). Yet, uncertainty about the situation and perceived exaggeration were associated with a reduced likeliness to implement the recommended protective behaviours during the 2009/10 pandemic (9). During the flu pandemic, a perceived inconsistency in recommendations was identified as a critical issue for non-compliance. Exaggeration of

risks often happens on social media, where especially highly emotional and often false information are shared (10). While a serial cross-sectional study involving over 13,000 participants during the 2009/2010 pandemic (11) showed that the internet was significantly less used as a source of information than traditional media, this may well have changed over the last decade. For example, the number of monthly Twitter users multiplied by ten from 30 million in 2009 to 330 million in 2019 (12) and Twitter seems to be seen as an alert tool in times of a crisis and a gateway for information (13). Thus, knowledge acquired during the last pandemic is only of limited value to guide crisis responses in the current outbreak.

The coronavirus is new, there is no vaccine or known effective treatment, case fatality rates are still uncertain. Psychologically, this means high uncertainty regarding the likelihood of catching the disease, its potential severity and ability to take control over the process by preventive measure. These perceptions are thus likely to be updated based on changes in epidemiology, media reports, information and misinformation.

As media and communication measures can influence these variables (7)(11) and as these are relevant for preparedness and protective behaviour (5)(14), the University of Erfurt collaboration aims at monitoring these variables during the current COVID-19 pandemic and to feed them into the communication process during the crisis. An additional aim is to reliably assess changes and shifts of risk perceptions and to identify the drivers and situations that are related to these shifts. How closely is risk perception related to actual risk? Further, it is important to understand the dynamics of risk perceptions, fears, misinformation and protective behaviours, understand which of the protective measures are known and which information is lacking. Based on this information it is possible to react to misinformation or suddenly increasing risk perceptions and panics.

## 5. References

1. Uscher-Pines L, Omer SB, Barnett DJ, Burke TA, Balicer RD. (2006) Priority setting for pandemic influenza: an analysis of national preparedness plans. *PLoS Med* 2006; **3**: 436.
2. Habersaat, K.B., Betsch, C., Danchin, M. et al. (2020) Ten considerations for effectively managing the COVID-19 transition. *Nat Hum Behav.* <https://doi.org/10.1038/s41562-020-0906-x>
3. Glik, D. C. (2007). Risk communication for public health emergencies. *Annual Review of Public Health*, 28, 33–54. <https://doi.org/10.1146/annurev.publhealth.28.021406.144123>
4. World Health Organization. (2017). Communicating risk in public health emergencies: A WHO guideline for emergency risk communication (ERC) policy and practice. World Health Organization.
5. Reynolds, B., & W. Seeger, M. (2005). Crisis and Emergency Risk Communication as an Integrative Model. *Journal of Health Communication*, 10(1), 43–55. <https://doi.org/10.1080/10810730590904571>
6. Rasmussen, S. A., & Goodman, R. A. (2018). *The CDC Field Epidemiology Manual*. Oxford University Press.
7. Reintjes R, Das E, Klemm C, Richardus JH, Keßler V, Ahmad A. “Pandemic Public Health Paradox”: Time Series Analysis of the 2009/10 Influenza A / H1N1 Epidemiology, Media Attention, Risk Perception and Public Reactions in 5 European Countries. *PLOS ONE* 2016; **11**: e0151258.
8. Van der Pligt J. Risk perception and self-protective behavior. *European Psychologist* 1996; **1**: 34–43.

9. Rubin, G. J., Amlot, R., Page, L., & Wessely, S. (2009). Public perceptions, anxiety, and behaviour change in relation to the swine flu outbreak: Cross sectional telephone survey. *BMJ*, 339(jul02 3), b2651–b2651. <https://doi.org/10.1136/bmj.b2651>
10. Vosoughi, S., Roy, D., & Aral, S. (2018). The spread of true and false news online. *Science*, 359(6380), 1146– 1151. <https://doi.org/10.1126/science.aap9559>
11. Walter, D., Böhmer, M. M., Reiter, S., Krause, G., & Wichmann, O. (2012). Risk perception and informationseeking behaviour during the 2009/10 influenza A(H1N1)pdm09 pandemic in Germany. *Eurosurveillance*, 17(13), 20131.
12. Statista. (2019). Number of monthly active Twitter users worldwide from 1st quarter 2010 to 1st quarter 2019.
13. Eriksson, M., & Olsson, E.-K. (2016). Facebook and Twitter in Crisis Communication: A Comparative Study of
14. Carpenter, C. J. (2010). A meta-analysis of the effectiveness of health belief model variables in predicting behavior. *Health Communication*, 25(8), 661–669. <https://doi.org/10.1080/10410236.2010.521906>
15. Betsch C, Bach Habersaat K, Deshevoi S, et al. Sample study protocol for adapting and/or translating the 5C scale to assess the psychological antecedents of vaccination. *BMJ Open* 2020;0:e034869. doi:10.1136/bmjopen-2019-034869.
16. Betsch C, Wieler L, Habersaat K, and the COSMO consortium (submitted). Rapid and flexible monitoring tool for behavioural insights related to COVID-19 across countries.
17. Repository for frameworks: [http://www.ncchpp.ca/708/repertoire\\_of\\_frameworks.ccnpps](http://www.ncchpp.ca/708/repertoire_of_frameworks.ccnpps) Including a framework on ethics during epidemics: <https://neac.health.govt.nz/system/files/documents/publications/getting-together-jul07.pdf>.
18. Nuffield Council of Bioethics: <https://www.nuffieldbioethics.org/publications/research-in-global-health-emergencies>
19. Taherdoost, H. (2017). Determining sample size; how to calculate survey sample size. *International Journal of Economics and Management Systems*, 2.

## Annex 1: Questionnaire

The following is the recommended standard approach. The most current version of the standard approach can be shared by WHO as an adaptable word version. Please contact Katrine Habersaat ([habersaatk@who.int](mailto:habersaatk@who.int)) or Martha Scherzer ([scherzerm@who.int](mailto:scherzerm@who.int)).

The situation evolves, sometimes fast and unevenly across the world, and so tailoring of the questionnaire to specific contexts is critical. Each country can select which variables they wish to focus on and can adapt items to their specific needs. Words highlighted in **yellow** indicate that context adaptation will be required.

Please note the text of the Introduction and other instructions is designed for online participants. If you conduct interviews by telephone this language should be revised accordingly.

Page 1	Dear Participant,
Introduction	<p>Thank you for taking part in this study to help improve actions taken in response to COVID-19 pandemic and to inform the response to similar future outbreaks.</p> <p>This study will involve answering a <b>XX</b>-minute survey which will be asking you questions relating to the coronavirus. Please do not start until you will have enough time to complete it in one go. Please close other programmes (e.g. chat or e-mail) to avoid distractions.</p> <p>This study is conducted by <b>[insert name of controller]</b> and the data is collected by <b>[insert name of the data collection agency]</b>.</p> <p>By taking part, you are agreeing that you have read and understood the information about the study below. Please ensure you have read and understood this information before continuing.</p> <p><b>What is this project about, and do I have to take part?</b></p> <p>This study aims to inform governmental outbreak response measures, including policies, interventions and communications. The information collected through this survey is important to support the implementation of specific programmatic interventions and policies in addition to the messaging necessary to encourage uptake of those measures. Participation is open to people at the age of 18 or over, living in <b>[insert name of country]</b> and is entirely voluntary. You do not have to be in isolation to take part.</p> <p><b>What are the benefits and risks of taking part?</b></p> <p>You may benefit from taking part in the survey by being motivated to look up information about the coronavirus pandemic. We will provide you with good resources at the end of the study. There are no foreseeable risks for you when taking part in the survey</p>



other than time spent on the survey and potential discomfort. Should you feel uncomfortable and want to leave the study you are free to do so without any consequences.

**What will you ask and what will happen to the information I give you?**

You will be asked questions about yourself, your knowledge of the coronavirus, the actions you have taken to protect yourself from the virus, your trust in various stakeholders, and your own fears and worries relating to the coronavirus pandemic. Some of these questions are considered sensitive data, such as questions relating to your trust in your government. However, you will not be asked to provide any personal data. Your anonymous data will be collected by [insert name of the data collection agency] and analysed by [insert name of controller]. Your data will be shared, but only with relevant researchers and government agencies. However, your data will be completely anonymous, and it will not be possible to identify you individually from your answers. This study has received approval from the WHO Research Ethics Review Committee and [insert name of relevant national or university ethics review committee].

**How long will my data be stored for?**

In order to help inform future pandemic and epidemic preparedness, the data you have provided will be helpful even beyond the current coronavirus pandemic. Your anonymous data will therefore be stored securely for up to 10 years by [insert name of controller] after the end of the research for this study. At this point the data will be reviewed, and if they are still deemed to be of public interest, they may be retained for longer. If not, your data will be permanently deleted.

**Local Data Protection Privacy Notice**

Notice: The controller for this project will be [insert name of controller]. The data will be collected by [insert name of the data collection agency].

This 'local' privacy notice sets out the information that applies to this particular study. Further information on how [insert name of controller] uses participant information can be found in the 'general' privacy notice: [provide relevant link to general data protection information of the controller].

The information that is required to be provided to participants under data protection legislation (GDPR [and/or other national data protection laws]) is provided across both the 'local' and 'general' privacy notices.

The lawful bases used in this survey are that it is undertaken as a task in the public interest and necessary for research and public health purposes, in accordance with the General Data Protection Regulation and national laws.

**Concerns**

If you are concerned about this study, or how your data is being processed, or if you would like to contact us about your rights, please get in touch with [insert name of controller] in the first instance at [insert contact email address].

**Consent**

I understand that:

- My participation is completely voluntary.

	<ul style="list-style-type: none"> <li>• All my answers will be used for scientific research to improve actions taken in response to the coronavirus pandemic and to inform the response to similar future outbreaks.</li> <li>• My data will be stored securely, however, no personal data will be stored, and my answer will be completely anonymous.</li> <li>• My data gathered in this study will be shared with relevant researchers and government agencies.</li> <li>• Because I am submitting anonymous data, it will not be possible to withdraw my answers after they have been submitted.</li> </ul> <p>Please note that you can stop the survey at any time. This will <u>not</u> entail any penalty, and it will <u>not</u> affect the services (health care services or others) that you receive.</p> <p>By ticking the box, you are agreeing that you are at least 18 years old, that you have read the information about the study, and that you voluntarily agree to take part in it.</p> <p><input type="checkbox"/> I agree to participate in this study.</p>
<p>Page 2</p> <p>Variable: socio-demography</p> <p>[Screen out: &lt;18]</p> <p>[Not in random order]</p>	<p><b>How old are you?</b></p> <p>I am ____ years old.</p> <p><b>What is your sex?</b></p> <p><input type="checkbox"/> Male</p> <p><input type="checkbox"/> Female</p> <p><input type="checkbox"/> Other</p> <p><b>How many years of education have you completed?</b></p> <p><b>Adapt to local context</b></p> <p><input type="checkbox"/> 0-9 years</p> <p><input type="checkbox"/> 10-12 years (secondary school completed)</p> <p><input type="checkbox"/> more than 12 years</p> <p><b>Are you a health professional?</b></p> <p><input type="checkbox"/> No</p> <p><input type="checkbox"/> Yes</p>

**Do you have a chronic illness?**

Yes

No

Don't know

**Where do you live?**

Rural area

Urban area

**In which district do you live?**

**[Dropdown list with all regions of country]**

**Who lives in your household besides yourself?**

*Choose as many as apply*

I live alone

I live with children under 18

I live with people in a COVID-19 risk group (people over 65 years and/or with chronic disease)

None of the above

**Please assess your private financial situation over the past three months:**

Improved

Remains the same

Worse

Don't know

Please click CONTINUE to proceed

<p>Page 3</p> <p>Variable: COVID-19 personal experience</p> <p>[Single choice]</p> <p>[Multiple choice for yes]</p> <p>[not in random order]</p>	<p><b>To your knowledge, are you, or have you been, infected with COVID-19?</b></p> <p>[*] No</p> <p>[*] Yes</p> <p>If “yes”:</p> <p><b>Was it:</b></p> <p>[*] Mild</p> <p>[*] Severe</p> <p><b>Was it:</b></p> <p>[*] Confirmed by a test</p> <p>[*] Not confirmed by a test</p> <p><b>Do you know people in your immediate social environment who are or have been infected with COVID-19 (suspected or confirmed)?</b></p> <p>[*] No</p> <p>[*] Yes</p> <p>If “yes”:</p> <p><b>Do you know someone who died from COVID-19?</b></p> <p>[*] No</p> <p>[*] Yes</p> <p>Please click CONTINUE to proceed</p>
<p>Page 4</p> <p>Variable: Health literacy</p> <p>[Not in random order]</p>	<p><b>How easy or difficult would you say it is to...:</b></p> <p><b>...find the information <u>you need</u> related to COVID-19?</b> [Very difficult [*][*][*][*][*][*][*] Very easy]</p> <p><b>...understand information about what to do if you think you have COVID-19?</b> [Answer scheme, see above]</p> <p><b>...judge if the information about COVID-19 in the media is reliable?</b> [Answer scheme, see above]</p>

	<p>...understand restrictions and recommendations of authorities regarding COVID-19? [Answer scheme, see above]</p> <p>...follow the recommendations on how to protect yourself from COVID-19? [Answer scheme, see above]</p> <p>...understand recommendations about when to stay at home from work/school, and when not to? [Answer scheme, see above]</p> <p>...follow recommendations about when to stay at home from work/school, and when not to? [Answer scheme, see above]</p> <p>...understand recommendations about when to engage in social activities, and when not to? [Answer scheme, see above]</p> <p>...follow recommendations about when to engage in social activities, and when not to? [Answer scheme, see above]</p> <p>Please click CONTINUE to proceed</p>
<p>Page 5</p> <p>Variable: Probability and Severity</p> <p>[random order of items]</p>	<p><b>What do you consider to be your own probability of getting infected with COVID-19?</b> Extremely unlikely [*] [*] [*] [*] [*] [*] [*] [*] Extremely likely</p> <p><b>How susceptible do you consider yourself to an infection with COVID-19?</b> Not at all susceptible [*] [*] [*] [*] [*] [*] [*] Very susceptible</p> <p><b>How severe would contracting COVID-19 be for you (how seriously ill do you think you will be)?</b> Not severe [*] [*] [*] [*] [*] [*] [*] Very severe</p> <p>Please click CONTINUE to proceed</p>
<p>Page 6</p> <p>Variable: Preparedness and Perceived self-efficacy</p> <p>[random order of items]</p>	<p><b>Next, we would like to know about you own practices related to COVID-19.</b></p> <p><b>I know how to protect myself from coronavirus</b> Not at all [*] [*] [*] [*] [*] [*] [*] Very much so</p> <p><b>For me avoiding an infection with COVID-19 in the current situation is...</b> Extremely difficult [*] [*] [*] [*] [*] [*] [*] Extremely easy</p> <p>Please click CONTINUE to proceed</p>
<p>Page 7</p> <p>Variable: Prevention – own behaviours</p>	<p><b>During the last 7 days, which of the following measures have you taken to prevent infection from COVID-19?</b></p> <p><b>Country adaptation: Adjust to national recommendations (include potentially widespread misperceptions as well as the official recommendations)</b></p> <p><i>Choose as many as apply</i></p> <p><b>Frequently washed my hands with soap and water for at least 20 seconds</b> Not at all [*] [*] [*] [*] [*] [*] [*] Very much so / Not a pp [*]</p>

<p>[random order of items]</p>	<p><b>Avoided touching my eyes, nose and mouth with unwashed hands</b> [Answer scheme, see above]</p> <p><b>Used disinfectants to clean hands when soap and water were not available</b> [Answer scheme, see above]</p> <p><b>Avoided a social event I wanted to attend</b> [Answer scheme, see above]</p> <p><b>Stayed at home from work/school</b> [Answer scheme, see above]</p> <p><b>Used antibiotics to prevent or treat COVID-19</b> [Answer scheme, see above]</p> <p><b>Wore a mask in public</b> [Answer scheme, see above]</p> <p><b>Ensured physical distancing in public</b> [Answer scheme, see above]</p> <p><b>Disinfected surfaces</b> [Answer scheme: see "Hand washing"]</p> <p style="text-align: center;">Please click CONTINUE to proceed</p>
<p>Page 8</p> <p>Variable: Affect</p> <p>[Random order of items]</p>	<p><b>Please choose one option per row below. COVID-19 to me feels ...</b></p> <p style="text-align: center;">close to me [*][*][*][*][*][*][*] far away from me</p> <p style="text-align: center;">Spreading slowly [*][*][*][*][*][*][*] Spreading fast</p> <p style="text-align: center;">Something I think about all the time [*][*][*][*][*][*][*] Something I almost never think about</p> <p style="text-align: center;">Fear-inducing [*][*][*][*][*][*][*] Not fear-inducing</p> <p style="text-align: center;">Media hyped [*][*][*][*][*][*][*] Not media hyped</p> <p style="text-align: center;">Something that makes me feel helpless [*][*][*][*][*][*][*] Something I am able to combat with my own action</p> <p style="text-align: center;">Stressful [*][*][*][*][*][*][*] Not stressful</p> <p style="text-align: center;">Please click CONTINUE to proceed</p>
<p>Page 9</p> <p>Variable: Trust in sources of information</p> <p>[Random order of items]</p>	<p><b>How much do you trust information about COVID-19 from the following sources?</b></p> <p><b>Television</b> Very little trust [*][*][*][*][*][*][*] A great deal of trust</p> <p><b>Newspapers</b> [Answer scheme, see above]</p> <p><b>Health workers</b> [Answer scheme, see above]</p>

	<p><b>Social media</b> [Answer scheme, see above]</p> <p><b>Radio</b> [Answer scheme, see above]</p> <p><b>Ministry of Health</b> [Answer scheme, see above]</p> <p><b>Institute of Public Health/Center for Disease Control</b> [Answer scheme, see above]</p> <p><b>Celebrities and social media influencers</b> [Answer scheme, see above]</p> <p><b>World Health Organization (WHO)</b> [Answer scheme, see above]</p> <p><b>COVID-19 Hotlines</b> [Answer scheme, see above]</p> <p><b>National COVID-19 information website</b> [Answer scheme, see above]</p> <p style="text-align: right;">Please click CONTINUE to proceed</p>
<p>Page 10</p> <p>Variable: Use of sources of information</p> <p>[Random order of items]</p>	<p><b>How often do you use the following sources for information about COVID-19?</b></p> <p><b>Television</b> Never [*] [*] [*] [*] [*] [*] [*] Very often</p> <p><b>Newspapers</b> [Answer scheme, see above]</p> <p><b>Health workers</b> [Answer scheme, see above]</p> <p><b>Social media</b> [Answer scheme, see above]</p> <p><b>Radio stations</b> [Answer scheme, see above]</p> <p><b>Ministry of Health</b> [Answer scheme, see above]</p> <p><b>Institute of Public Health</b> [Answer scheme, see above]</p> <p><b>Celebrities and social media influencers</b> [Answer scheme, see above]</p> <p><b>World Health Organization (WHO)</b> [Answer scheme, see above]</p> <p><b>COVID-19 Hotlines</b> [Answer scheme, see above]</p> <p><b>National COVID-19 information website</b> [Answer scheme, see above]</p> <p style="text-align: right;">Please click CONTINUE to proceed</p>

<p>Page 11</p> <p>Variable: Frequency of Information</p>	<p><b>How often do you seek information about COVID-19?</b></p> <p>Never [*] [*] [*] [*] [*] [*] [*] Several times a day</p> <p>Please click CONTINUE to proceed</p>
<p>Page 12</p> <p>Variable: Trust in institutions (perceptions)</p> <p>[Random order of items]</p>	<p><b>How much confidence do you have that the following can handle the COVID-19 challenge well?</b></p> <p><b>Your family doctor</b> Very low confidence [*] [*] [*] [*] [*] [*] [*] Very high confidence / Not applicable [*]</p> <p><b>Your employer</b> [Answer scheme, see above]</p> <p><b>Hospitals</b> [Answer scheme, see above]</p> <p><b>Ministry of Health</b> [Answer scheme, see above]</p> <p><b>Institute of Public Health/Center for disease Control</b> [Answer scheme, see above]</p> <p><b>Schools</b> [Answer scheme, see above]</p> <p><b>Public transportation companies</b> [Answer scheme, see above]</p> <p><b>Police</b> [Answer scheme, see above]</p> <p><b>Your church/place of worship</b> [Answer scheme, see above]</p> <p>Please click CONTINUE to proceed</p>
<p>Page 14</p> <p>Variable: Conspiracies (perceptions)</p> <p>[random order of items]</p>	<p><b>Please consider the decisions that are made in your country to reduce spread of COVID-19:</b></p> <p><b>I think that....</b></p> <p>...many very important things happen in the world, which the public is never informed about certainly not true [*] [*] [*] [*] [*] [*] [*] certainly true</p> <p>... politicians usually do not tell us the true motives for their decisions [See answer scheme above.]</p> <p>... government agencies closely monitor all citizens [See answer scheme above.]</p> <p>... events which superficially seem to lack a connection are often the result of secret activities [See answer scheme above.]</p> <p>... there are secret organizations that greatly influence political decisions [See answer scheme above.]</p> <p>Please click CONTINUE to proceed</p>
<p>Page 15</p>	<p><b>Please consider your experience during COVID-19 pandemic:</b></p>



<p>Variable: Resilience [random order of items]</p>	<p><b>I have a hard time making it through stressful events</b> Strongly disagree [*][*][*][*][*][*][*] Strongly agree</p> <p><b>It does not take me long to recover from a stressful event</b> [See answer scheme above.]</p> <p><b>It is hard for me to snap back when something bad happens</b> [See answer scheme above.]</p> <p>Please click CONTINUE to proceed</p>
<p>Page 16 Variable: Testing and tracing  [Multiple choice for all four answer options]  [Overall two question not in random order]  [Answer options under each answer option on random order]</p>	<p><b>If you have been in contact with someone who tested positive for COVID-19 and have no symptoms yourself – will you get tested if you have the opportunity?</b></p> <p>[*] I would get tested for sure [leads to additional answer options below]</p> <p>[*] I may not get tested [leads to additional answer options below]</p> <p><u>For those who select “I would get tested for sure”:</u></p> <p><b>Please elaborate on this</b></p> <p><i>Choose as many as apply</i></p> <p><b>I would get tested for sure because...</b></p> <p>[*]... I want to receive the appropriate care in case of a positive test</p> <p>[*]...this is my responsibility as a citizen</p> <p>[*]...I would face penalties if I did not</p> <p>[*]...I believe this helps stop the spread of COVID-19</p> <p>[*]...this way I can protect other people</p> <p>[*]...my friends and family would expect me to get tested</p> <p><u>For those who select “I may not get tested”:</u></p> <p><b>Please elaborate on this</b></p> <p><i>Choose as many as apply</i></p> <p><b>I may not get tested because...</b></p> <p>[*]... getting tested would cost money (e.g. transportation, buying the test, taking time off work)</p>

[\*]... I do not know where to go to be tested

[\*]... it is too time-consuming to get tested

[\*]... this will result in loss of income for me due to quarantine while waiting to get the results

[\*]... this would result in loss of income for me if I get a positive test

[\*]... people might blame me for my actions if I get a positive test

[\*]... I might face fines or other penalties if I had violated official COVID restrictions

[\*]... I do not trust authorities with my personal data

[\*]... I do not believe COVID-19 exists

[\*]... there is nothing I can do, even if I get a positive test

[\*]... I am not able to self-isolate in case I get a positive test

[\*]... I do not think the tests are reliable

[\*]... I am worried people will treat me badly if I get a positive test

[\*]... I am worried I will get infected at the testing site

[\*]... I think testing will be painful

**If you test positive for COVID-19 and are asked to share with health authorities the names of people you have been in contact with – will you share all names?**

[\*] I would share all names for sure [leads to additional answer options below]

[\*] I may not share all names [leads to additional answer options below]

For those who select “I would get tested for sure”:

**Please elaborate on this**

*Choose as many as apply*

**I would share all names for sure because...**

	<p>[*]... I believe this helps stop spread of COVID-19</p> <p>[*]... this is my responsibility as a citizen</p> <p>[*]... this way I can protect other people</p> <p>[*]... my friends and family would expect me to do this</p> <p>[*]... I would face penalties if I did not</p> <p><u>For those who select "I may not share all names"</u></p> <p><b>Please elaborate on this</b></p> <p><i>Choose as many as apply</i></p> <p><b>I may not share all names because...</b></p> <p>[*]... I could contact them myself</p> <p>[*]... I believe this could result in loss of income for those people due to quarantine</p> <p>[*]... I believe people would blame me for having shared their name</p> <p>[*]... I do not trust authorities</p> <p>[*]... my family and friends would expect me not to share names</p> <p>[*]... I would cause inconvenience for the people whose names I share</p> <p>[*]... I do not want others to know I tested positive</p> <p>Please click CONTINUE to proceed</p>
<p>Page 17</p> <p>Variable: Fairness</p> <p>[random order of items]</p>	<p><b>Please consider the decisions that are made in your country to reduce spread of COVID-19:</b></p> <p><b>I think the decisions are fair</b> Strongly disagree [*] [*] [*] [*] [*] [*] [*] Strongly agree</p> <p><b>I would convince others that the decisions are right</b> Strongly disagree [*] [*] [*] [*] [*] [*] [*] Strongly agree</p> <p>Please click CONTINUE to proceed</p>
<p>Page 18</p>	<p><b>Please now give your opinion on the following statements:</b></p>

<p>Lifting restrictions (pandemic transition phase)</p> <p>[random order of items]</p>	<p><b>If a COVID-19 vaccine becomes available and is recommended for me, I would get it.</b> Strongly disagree [*] [*] [*] [*] [*] [*] [*] Strongly agree</p> <p><b>In the event of an outbreak it's appropriate to avoid certain people on the basis of their ethnicity.</b> Completely disagree [*] [*] [*] [*] [*] [*] [*] Completely agree</p> <p><b>I think that the restrictions currently being implemented are greatly exaggerated.</b> [See answer scheme above.]</p> <p><b>The government should be allowed to force people into self-isolation if they have been in contact with someone who was infected</b> [See answer scheme above.]</p> <p><b>More tests for coronavirus infection should be carried out in the population</b> [See answer scheme above.]</p> <p><b>I am worried that the pandemic will have economic consequences for me in the future</b> [See answer scheme above.]</p> <p><b>Some restrictions have now been changed related to COVID-19. Please indicate, to which degree you support the following decisions:</b></p> <p><b>Country adaptation: Add 3-5 decisions that were implemented. Examples include:</b></p> <p><b>Example: Compulsory face masks in closed public spaces</b> Strongly support [*] [*] [*] [*] [*] [*] [*] Do not support at all</p> <p><b>Example: Restricting restaurants to outside spaces</b> [See answer scheme above.]</p> <p><b>Example: Reopening of schools</b> [See answer scheme above.]</p> <p><b>There is a lot of debate, and some suggestions have been made regarding future restrictions related to COVID-19. What is your level of agreement with the following scenarios:</b></p> <p><b>Country adaptation: Add 3-5 decisions that are being considered or discussed in the public debate. Examples include:</b></p> <p><b>Example: Ban on mass gatherings in streets</b> Strongly support [*] [*] [*] [*] [*] [*] [*] Do not support at all</p> <p><b>Example: Introduction of mandatory testing of school teachers</b> [See answer scheme above.]</p> <p><b>Example: Opening borders to more countries</b> [See answer scheme above.]</p> <p>Please click CONTINUE to proceed</p>
<p>Page 19</p>	<p><b>Within the last 2 weeks, have you done the following...?</b></p>

<p>Variable: Unwanted behaviour</p> <p>[Integrate in randomized order of the policy items]</p> <p>[Random order of items]</p>	<p>[*] Yes [*] No [*] Not applicable</p> <p><b>Avoided people that I thought might infect me, based on their ethnicity</b> [See answer scheme above.]</p> <p><b>Exercised less than I did before the pandemic</b> [See answer scheme above.]</p> <p><b>Drank more alcohol than I did before the pandemic</b> [See answer scheme above.]</p> <p><b>Ate more unhealthy food than I did before the pandemic</b> [See answer scheme above.]</p> <p><b>Smoked more than I did before the pandemic</b> [See answer scheme above.]</p> <p><b>Postponed vaccination for myself or my child</b> [See answer scheme above.]</p> <p><b>Avoided going to the doctor for a non-COVID-19-related problem</b> [See answer scheme above.]</p> <p><b>Bought drugs that I heard are good for treating COVID-19</b> [See answer scheme above.]</p> <p>Please click CONTINUE to proceed</p>
<p>Page 20</p> <p>Variable: Wellbeing</p> <p>[Random order of items]</p>	<p><b>We would now like for you to indicate your general well-being:</b></p> <p>Over the past 2 weeks...</p> <p><b>... I have felt cheerful and in good spirits</b> [All of the time – Most of the time – More than half the time – Less than half the time – Some of the time – At no time]</p> <p><b>... I have felt calm and relaxed</b> [See answer scheme above.]</p> <p><b>... I have felt active and vigorous</b> [See answer scheme above.]</p> <p><b>... I woke up feeling fresh and rested</b> [See answer scheme above.]</p> <p><b>... my daily life has been filled with things that interest me</b> [See answer scheme above.]</p> <p>Please click CONTINUE to proceed</p>
<p>Page 21</p>	<p><b>There are currently no vaccines available to prevent COVID-19, but many researchers are working to develop and test vaccines. Please share your position on a potential future COVID-19 vaccine:</b></p>

<p>Variable: COVID-19 vaccine</p> <p>[First three items: random order]</p> <p>[Fourth item: answer options in random order]</p> <p>[Fifth item: always last]</p>	<p><b>I believe a vaccine can help control the spread of COVID-19</b> Strongly disagree [*] [*] [*] [*] [*] [*] Strongly agree</p> <p><b>If I knew I had been infected with COVID-19 before, I would not get the vaccine even if it were available</b> [See answer scheme above.]</p> <p><b>When everyone else is vaccinated against COVID-19, then I don't have to get vaccinated</b> [See answer scheme above.]</p> <p><b>If a COVID-19 vaccine is made available in my country, my decision of whether or not to get vaccinated would depend on:</b>  <i>Choose as many as apply</i></p> <p>Country in which the vaccine is produced Not at all [*] [*] [*] [*] [*] [*] Very much so</p> <p>Recommendation from my family doctor [See answer scheme above.]</p> <p>Recommendation of the Ministry of Health [See answer scheme above.]</p> <p>Whether the vaccine has been in use for a long time with no serious side-effects [See answer scheme above.]</p> <p>Whether the vaccine is used in other countries [See answer scheme above.]</p> <p>Risk of getting infected with COVID-19 at the time when the vaccine is available [See answer scheme above.]</p> <p>How easy it is to get the vaccine (e.g. available out-of-hours or in pharmacies) [See answer scheme above.]</p> <p>Whether the vaccine is free of charge [See answer scheme above.]</p> <p>Whether a high vaccination uptake would lift restrictions on movement and gathering in groups [See answer scheme above.]</p> <p><b>Apart from COVID-19, I think everyone should be vaccinated according to the national vaccination schedule</b></p> <p>[*] Yes</p> <p>[*] No</p> <p>[*] Don't know</p> <p>Please click CONTINUE to proceed</p>
<p>Page 22</p>	<p>Debriefing</p>

Debriefing	<p>Thank you very much!</p> <p>Your participation provides valuable insights for all of us to react appropriately in the current COVID-19 situation and to reach all citizens with useful information in a timely manner.</p> <p>For information about COVID-19, please visit the following websites: <b>Adapted to national setting: add one or more trusted information sources</b></p> <p>If you have any questions, please contact <b>Adapted to national setting: add contact person</b> in <b>implementing country</b>.</p> <p>If you have changed your opinion and would like to withdraw your consent to use your data, please click on "Withdraw my consent". This will <u>not</u> entail any penalty, and it will <u>not</u> affect the services (health care services or others) that you receive.</p> <p><input type="checkbox"/> Withdraw my consent</p> <p>Please click CONTINUE to finish the survey</p>
------------	---

## Annex 2: Data analysis and presentation of results

Data can be analysed with any data analysis software.

In order to facilitate the process of analysing and presenting the data, we offer a RMarkdown notebook (based on the free software R). These notebooks blend analysis code with result output like plots and regression tables as well as explanatory text. The University of Erfurt collaboration group implemented a first template that is offered as a blueprint for other countries.

Please note:

As the questionnaire is adapted to country specific features (e.g. different states and authorities) the notebook has to be tailored as well. Consequently, before collecting the first data, a custom notebook needs to be created for each subgroup.

Each notebook is hosted on RStudio cloud, a free to use online platform where questionnaire data can be uploaded. With a single click, notebooks can be executed then, i.e. data analyses are run and a website is generated visualizing the main results (see Figure 3).

The code is offered as a \*.zip file at <https://hdl.handle.net/20.500.12034/2392> and can be uploaded to a RStudio cloud.

Please check for updates in the repository (<https://hdl.handle.net/20.500.12034/2392>) where new examples of code will be offered. We suggest that countries also share their code as part of survey documents using open source research platforms, allowing other countries to use the code on their data.

It is important that the data set uses the variable labels and codes as provided in Annex 1 to allow the script to run without errors.



Figure 3: Data analysis workflow using RMarkdown notebooks

The image shows a screenshot of the RStudio Cloud interface. The main window displays an R Markdown notebook with R code for data loading and variable definition. The code includes comments and R commands for reading Excel files, binding rows, and defining factors for age groups and gender. The environment pane on the right shows the loaded data files: 'data' (1946 obs. of 251 variables), 'data\_freq' (969 obs. of 241 variables), 'data\_info...' (3 obs. of 3 variables), and 'data\_preve...' (48 obs. of 4 variables). The file pane shows the project structure with files like 'analyses.html', 'analyses.Rmd', 'analyses\_files', 'custom.css', 'data.t1.xlsx', and 'data.t2.xlsx'. A callout box labeled '1 Upload your data' points to the file pane. A second callout box labeled '2 Run the notebook' points to the 'Run' button in the R Markdown editor. A third callout box labeled '3 Result website is created' points to a browser window showing the rendered HTML report. The report contains a table of contents with items 1 through 10, where item 6 'Confidence in crisis management' is highlighted. Below the table of contents are three charts: 'Trust in media' (a horizontal bar chart showing mean values for various media types), 'Relevance of media' (a horizontal bar chart showing percentages for different media types), and '6 Confidence in crisis management' (a vertical bar chart showing trust levels for different health sector entities).

**1 Upload your data**

**2 Run the notebook**

**3 Result website is created**

**6 Confidence in crisis management**

**Trust in media**  
Results from recent survey (18/03/2020)  
Rated on scales ranging from 1 (not at all) to 7 (very much).  
Mean values and 95% confidence intervals.

Media Type	Mean Value
Public TV	4.05
Public radio	4.26
Conversations	4.07
Newspapers	4.12
Magazines	3.98
Private radio	3.87
Search engines	3.81
Private TV	3.98
News websites	3.54
Social media	2.81

**Relevance of media**  
Results from recent survey (18/03/2020)  
Calculated as the product of usage frequency and trust.  
Mean values and 95% confidence intervals.

Media Type	Percentage
Conversations	46%
Public TV	44%
Public radio	39%
Search engines	37%
Newspapers	34%
Private TV	31%
News websites	27%
Private radio	27%
Social media	26%
Magazines	25%

**6 Confidence in crisis management**  
**Trust in crisis management (1): Health sector**  
Rated on scales ranging from 1 (not at all) to 7 (very much trust).  
Mean values and 95% confidence intervals.

Entity	Mean Value
Own doctor	5.14
Family	4.81
Local health department	4.73
Medical association	4.69
Healthcare provider	4.60
Local health department	4.54
Medical association	4.50

### **The WHO Regional Office for Europe**

The World Health Organization (WHO) is a specialized agency of the United Nations created in 1948 with the primary responsibility for international health matters and public health. The WHO Regional Office for Europe is one of six regional offices throughout the world, each with its own programme geared to the particular health conditions of the countries it serves.

#### **Member States**

Albania  
Andorra  
Armenia  
Austria  
Azerbaijan  
Belarus  
Belgium  
Bosnia and Herzegovina  
Bulgaria  
Croatia  
Cyprus  
Czechia  
Denmark  
Estonia  
Finland  
France  
Georgia  
Germany  
Greece  
Hungary  
Iceland  
Ireland  
Israel  
Italy  
Kazakhstan  
Kyrgyzstan  
Latvia  
Lithuania  
Luxembourg  
Malta  
Monaco  
Montenegro  
Netherlands  
North Macedonia  
Norway  
Poland  
Portugal  
Republic of Moldova  
Romania  
Russian Federation  
San Marino  
Serbia  
Slovakia  
Slovenia  
Spain  
Sweden  
Switzerland  
Tajikistan  
Turkey  
Turkmenistan  
Ukraine  
United Kingdom  
Uzbekistan

### **World Health Organization Regional Office for Europe**

UN City, Marmorvej 51,  
DK-2100 Copenhagen Ø, Denmark  
Tel.: +45 45 33 70 00 Fax: +45 45 33 70 01  
Email: [eurocontact@who.int](mailto:eurocontact@who.int)  
Website: [www.euro.who.int](http://www.euro.who.int)